

Patient Details

Forename

Surname

Protocol

Page:1 of 1

DOB

Patient NO

Local No.

Course Name:

Consultant

Ward

Type of line
No. of lumen:

Trastuzumab (SC) 600mg 3 weekly

Diagnosis

Carcinoma of Breast

NHS No

[illegible]

Additional Prescribing Notes

Regime emetogenicity: NOT emetogenic.



MUGA scan required prior to day 1 and then every 4 months during treatment.

Administer as a SC injection over 2-5 minutes.
Alternated the injection site between the right and left thigh. New injections should be given at least 2.5cm from the old site and never into areas where the skin is red, bruised, tender or hard.

FIRST INJECTION: Observed for six hours after the first injection

SECOND INJECTION: Observe for two hours after second injection

SUBSEQUENT INJECTIONS: If no administration-related reactions are seen with previous injections patients do not need to be observed.

| Day | Date and Time | Drug and dose (per m2) or dose (per kg) | ACTUAL DOSE | Infusion Fluid and Final Volume | Route | Additives | Time/Infusion Rate | Line | Given/Checked by | Time Start/Stop | Comments |
|-----|---------------|---|---------------|---------------------------------|-------|-----------|--------------------|------|--|---|-----------------------------|
| 1 | T=:hrs | TRASTUZUMAB (SUB CUT) (600mg) | 600 mg | None | SC | | Slow Bolus | |  Batch No. |  | Administer over 2-5 minutes |

| | | | | |
|----------------------|----------------------|-----------------------|---------------------------------|---|
| Allocated by: | Confirmed by: | Authorised by: | Checked by: (Pharmacist) | <div> <div>Parenteral</div> <div>Intrathecal</div> <div>Oral</div> </div> <div> <div>1</div> <div>0</div> <div>0</div> </div> |
| Date: | Date: | Date: | Date: | <div>Chart Id.:</div> |