

Patient Details

Forename \_\_\_\_\_ Surname \_\_\_\_\_

Address \_\_\_\_\_

DOB \_\_\_\_\_ Patient NO \_\_\_\_\_ Local No. \_\_\_\_\_

Consultant \_\_\_\_\_ Ward \_\_\_\_\_

NHS No \_\_\_\_\_

Protocol

Escalated BEACOPDAC

Course Name: Escalated BEACOPDac SA<2.1m2 cycle 2+

Type of line \_\_\_\_\_ No. of lumen: \_\_\_\_\_

Diagnosis

SA (m²)  
Height (m)  
Weight (kg)

Page:1 of 7

Monitoring	Acceptable Range		Date Due	Date of Test	Value	Checked
Height (m)						
Weight (kg)						
SA (m²)						
ALA TRANSAM 1.5ULN	0.00	60.00	Day 1			
BILIRUBIN 1.5ULN	0.00	31.50	Day 1			
COCKCROFT (>60)	60.00	300.00	Day 1			
NEUTROPHILS > 1.0	1.00	15.00	Day 1			
PLATELETS > 80	80.00	600.00	Day 1			
WHITE BLOOD CELL > 2.5	2.50	11.00	Day 1			

Additional Prescribing Notes

Doxorubicin and vincristine can cause pain and tissue necrosis if extravasated.

Doses capped at 2.1m/2.

Day	Date and Time	Drug and dose (per m2) or dose (per kg)	ACTUAL DOSE	Infusion Fluid and Final Volume	Route	Additives	Time/Infusion Rate	Line	Given/ Checked by	Time Start/ Stop	Comments
1	T=hrs	FREE FLOWING INFUSION (500ml)	500 ml	SODIUM CHLORIDE 0.9%	IV				<div></div> <div>Batch No.</div>	<div></div>	
1	T=hrs	AKYNZEO 300MG/0.5MG (1capsule)	1 capsule		PO				<div></div> <div>Batch No.</div>	<div></div>	To be given 60 minutes prior to chemotherapy
1	T=hrs	DOXORUBICIN (35mg/m²)	mg	None	IV		Slow Bolus		<div></div> <div>Batch No.</div>	<div></div>	
1	T=hrs	CYCLOPHOSPHAMIDE (1250mg/m²)	0 mg	SODIUM CHLORIDE 0.9%  250 ml	IV		Infuse over 30 Mins at a rate 500 ml/hr		<div></div> <div>Batch No.</div>	<div></div>	

Allocated by: \_\_\_\_\_ Date: \_\_\_\_\_

Confirmed by: \_\_\_\_\_ Date: \_\_\_\_\_

Authorised by: \_\_\_\_\_ Date: \_\_\_\_\_

Checked by: (Pharmacist) \_\_\_\_\_ Date: \_\_\_\_\_

Parenteral  
Intrathecal  
Oral

3  
0  
4

Patient Details

Forename

Surname

Protocol

Address

DOB

Patient NO

Local No.

Course Name:

NHS No

Ward

Escalated BEACOPDAC

Escalated BEACOPDac SA<2.1m2 cycle 2+

SA (m²)  
Height (m)  
Weight (kg)

Day	Date and Time	Drug and dose (per m2) or dose (per kg)	ACTUAL DOSE	Infusion Fluid and Final Volume	Route	Additives	Time/Infusion Rate	Line	Given/ Checked by	Time Start/ Stop	Comments
1	T=hrs	ETOPOSIDE (200mg/m²)	mg	SODIUM CHLORIDE 0.9% 1000 ml	IV		Infuse over 2 Hrs at a rate 500 ml/hr		<div></div> <div>Batch No.</div>	<div></div>	
2	T=hrs	FREE FLOWING INFUSION (500ml)	500 ml	SODIUM CHLORIDE 0.9%	IV				<div></div> <div>Batch No.</div>	<div></div>	
2	T=hrs	DACARBAZINE (250mg/m²)	mg	SODIUM CHLORIDE 0.9% 500 ml	IV		Infuse over 30 Mins at a rate 1000 ml/hr		<div></div> <div>Batch No.</div>	<div></div>	
2	T=hrs	ETOPOSIDE (200mg/m²)	mg	SODIUM CHLORIDE 0.9% 1000 ml	IV		Infuse over 2 Hrs at a rate 500 ml/hr		<div></div> <div>Batch No.</div>	<div></div>	
3	T=hrs	FREE FLOWING INFUSION (500ml)	500 ml	SODIUM CHLORIDE 0.9%	IV				<div></div> <div>Batch No.</div>	<div></div>	
3	T=hrs	DACARBAZINE (250mg/m²)	mg	SODIUM CHLORIDE 0.9% 500 ml	IV		Infuse over 30 Mins at a rate 1000 ml/hr		<div></div> <div>Batch No.</div>	<div></div>	
3	T=hrs	ETOPOSIDE (200mg/m²)	mg	SODIUM CHLORIDE 0.9% 1000 ml	IV		Infuse over 2 Hrs at a rate 500 ml/hr		<div></div> <div>Batch No.</div>	<div></div>	

Allocated by:	Confirmed by:	Authorised by:	Checked by: (Pharmacist)
Date:	Date:	Date:	Date:

Patient Details

ForenameSurnameProtocol

AddressDOBPatient NOLocal No.Course Name:

NHS No

Ward

Escalated BEACOPDAC

Escalated BEACOPDac SA<2.1m2 cycle 2+

SA (m²)  
Height (m)  
Weight (kg)

Page:3 of 7

Day	Date and Time	Drug and dose (per m2) or dose (per kg)	ACTUAL DOSE	Infusion Fluid and Final Volume	Route	Additives	Time/Infusion Rate	Line	Given/ Checked by	Time Start/ Stop	Comments
8	T=hrs	ONDANSETRON (8mg)	8 mg	None	PO				<div></div> <div>Batch No.</div>	<div></div>	
8	T=hrs	VINCRISTINE (1.4mg/m²)	mg	SODIUM CHLORIDE 0.9% 50 ml	IV				<div></div> <div>Batch No.</div>	<div></div>	Max dose: 2mg. Infuse over 5-10 minutes.
8	T=hrs	FREE FLOWING INFUSION (500ml)	500 ml	SODIUM CHLORIDE 0.9%	IV				<div></div> <div>Batch No.</div>	<div></div>	
8	T=hrs	HYDROCORTISONE (100mg)	100 mg	None	IV		Slow Bolus		<div></div> <div>Batch No.</div>	<div></div>	
8	T=hrs	BLEOMYCIN (10000unit/m²)	unit	None	IV		Slow Bolus		<div></div> <div>Batch No.</div>	<div></div>	

Allocated by:

Confirmed by:

Authorised by:

Checked by: (Pharmacist)

Date:

Date:

Date:

Date:

Patient Details

Forename	Surname		Protocol	ESCALATED BEACOPDAC										SA (m²)	
			Course Name	Escalated BEACOPDac SA<2.1m2 cycle 2+										Height (m)	
DOB	Patient NO		Local No.		NHS No								Weight (kg)		
Consultant			Ward		Diagnosis										
Address															

Record drug allergies or sensitivities

				Time	Date													
Drug & dose	PREDNISOLONE																	
Actual dose	mg		Duration	14 DAYS														
Route	PO		Start Date															
Frequency	OD		Start Day	1														
Quantity Dispensed		Dispensed by																
		Accuracy check																
Note																		
Drug & dose	METOCLOPRAMIDE																	
Actual dose	10 mg		Duration	PRN														
Route	PO		Start Date															
Frequency	TDS		Start Day	1														
Quantity Dispensed		Dispensed by																
		Accuracy check																
Note	If pre-pack supplied record Batch Number :_____.																	

Allocated by:	Confirmed by:	Authorised by:	Checked by: (Pharmacist)	
Date:	Date:	Date:	Date:	

Patient Details

Forename	Surname		Protocol	ESCALATED BEACOPDAC					SA (m²)				
			Course Name	Escalated BEACOPDac SA<2.1m2 cycle 2+					Height (m)				
DOB	Patient NO		Local No.		NHS No							Weight (kg)	
			Ward										
Address													

Record drug allergies or sensitivities

				Time	Date													
Drug & dose	CO-TRIMOXAZOLE																	
Actual dose	960 mg		Duration	21 DAYS														
Route	PO		Start Date															
Frequency	OD MWF		Start Day	1														
Quantity Dispensed		Dispensed by																
		Accuracy check																
Note	To be taken DAILY on Mondays, Wednesdays and Fridays. This is continuous treatment. If pre-pack supplied record Batch Number :_____.																	
Drug & dose	ACICLOVIR																	
Actual dose	200 mg		Duration	21 DAYS														
Route	PO		Start Date															
Frequency	TDS		Start Day	1														
Quantity Dispensed		Dispensed by																
		Accuracy check																
Note	This is continuous treatment supply original packs. If pre-pack supplied record Batch Number :_____.																	

Allocated by:	Confirmed by:	Authorised by:	Checked by: (Pharmacist)	
Date:	Date:	Date:	Date:	

Patient Details

Forename	Surname		Protocol	ESCALATED BEACOPDAC					SA (m²)				
			Course Name	Escalated BEACOPDac SA<2.1m2 cycle 2+					Height (m)				
DOB	Patient NO		Local No.		NHS No							Weight (kg)	
			Ward										
Address													

Record drug allergies or sensitivities

				Time	Date													
Drug & dose	RANITIDINE																	
Actual dose	150 mg		Duration	21 DAYS														
Route	PO		Start Date															
Frequency	BD		Start Day	1														
Quantity Dispensed			Dispensed by															
			Accuracy check															
Note	This is continuous treatment supply original packs. If pre-pack supplied record Batch Number : _____.																	
Drug & dose	METOCLOPRAMIDE																	
Actual dose	10 mg		Duration	PRN														
Route	PO		Start Date															
Frequency	TDS		Start Day	8														
Quantity Dispensed			Dispensed by															
			Accuracy check															
Note	Discuss with patient and delete if supply not required. If pre-pack supplied record Batch Number : _____.																	

Allocated by:	Confirmed by:	Authorised by:	Checked by: (Pharmacist)	
Date:	Date:	Date:	Date:	

Patient Details																								
Forename	Surname	Protocol	ESCALATED BEACOPDAC										SA (m²)											
		Course Name	Escalated BEACOPDac SA<2.1m2 cycle 2+										Height (m)											
DOB	Patient NO	Local No.	NHS No											Weight (kg)										
		Ward																						
Address																								
Record drug allergies or sensitivities																								
												Time	Date											
Drug & dose	FILGRASTIM (G-CSF)																							
Actual dose	microgram		Duration	5 DAYS																				
Route	SC		Start Date																					
Frequency	OD		Start Day	9																				
Quantity Dispensed		Dispensed by																						
		Accuracy check																						
Note	SUBCUTANEOUS BOLUS																							
	Discontinue when, after reaching nadir the WBC remains over 1.0 on 3 successive days																							

Allocated by:	Confirmed by:	Authorised by:	Checked by: (Pharmacist)	
Date:	Date:	Date:	Date:	