


# Optimal Vaccine Schedule in Immuno-compromised\* Patients

Based on current (November 2021) advice from the JCVI



\*Immuno-compromised includes: Chemotherapy or Radiotherapy or Prednisolone ≥ 7.5mg/day for 3 months, Prednisolone ≥ 10mg/day for 1 month in last 3 months or, Prednisolone ≥ 20mg/day for 10days in last month or Prednisolone ≥ 40mg in last month. First & second primary dose can be with Vaxzevria™ (0.5mL), Comirnaty™ (0.3mL) or Spikevax™ (0.5mL) - subject to other contra-indications. ‡Third primary dose should be Comirnaty™ (0.3mL) or Spikevax™ (0.5mL) unless contra-indicated, ideally in a treatment break but do not delay vaccination. †Booster dose should be Comirnaty™ (0.3mL) or Spikevax™ (0.25mL) unless contra-indicated, ideally during a treatment break or 2 weeks after completion of treatment (allow normal treatment cycle length plus 2 weeks), where this is not possible - in the few days immediately prior to next treatment when immune recovery is likely to be highest. Vaxzevria™ = AstraZeneca, Comirnaty™ = Pfizer, Spikevax™ = Moderna

Patients who have undergone Stem Cell Transplant should have all vaccinations repeated 6 months after transplantation